

SF00-SF06-0003

RAIDEN

Low Volume Abamectin Drench
For Sheep and Lambs



Active Content & Action

Abamectin 2.0 g/L. Abamectin is part of the Macrocylic Lactones group of chemicals. It works by interrupting the nervous systems of parasites, leading to paralysis and eventual death.

Use

For the treatment and control of abamectin sensitive strains of internal parasites (including benzimidazole, levamisole and morantel resistant strains) of sheep and itch mite and nasal bot of sheep including:

Adult and Immature Gastrointestinal Roundworms:

<i>Haemonchus contortus</i>	- Barbers pole worm (including inhibited L4 stage)
<i>Haemonchus placei</i>	- Large stomach worm
<i>Ostertagia circumcincta</i>	- Small brown stomach worms (including inhibited L4 stage)
<i>Trichostrongylus axei</i>	- Stomach hair worm
<i>Trichostrongylus colubriformis</i> ,	
<i>Trichostrongylus vitrinus</i>	- Black scour worm
<i>Cooperia curticei</i> , <i>Cooperia</i>	
<i>oncophora</i>	- Small intestinal worm
<i>Nematodirus spathiger</i> ,	
<i>Nematodirus filicollis</i>	- Thin necked intestinal worm
<i>Chabertia ovina</i>	- Large mouthed bowel worm
<i>Oesophagostomum columbianum</i>	- Nodule worm
<i>Oesophagostomum venulosum</i>	- Large bowel worm
<i>Trichuris ovis</i>	- Whipworm
<i>Strongyloides papillosus</i>	- Intestinal threadworm

Also Treats and Controls:

<i>Dictyocaulus viviparus</i>	- Large lungworm
<i>Psorergates ovis</i>	- Itch mite
<i>Oestrus ovis</i>	- Nasal bot

Indications

Nil

Information About Selenium:

Raiden does not contain selenium. Excess selenium can cause toxicity issues in sheep, particularly if they are suffering liver damage from eating particular weeds. Where sheep are receiving sufficient selenium in their diet, it may be better to use an unselenised drench.

Period of Protection

This is a short acting product and can only be expected to work where resistance is not prevalent. Always conduct drench resistance testing and seek veterinary advice if you are concerned for livestock wellbeing.

Doses

2000 doses per 10L pack
(average weight 50kg).

Pack Size(s) Available

10L

Australian Poisons Schedule

S5

Presentation

50 drums to a pallet

APVMA Approval No.:

86851



Dosage and Administration FOR EXTERNAL USE ONLY.

Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs). Do not underdose.

A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Safe to use on pregnant and non-pregnant animals of any age when used as directed.

Dose sheep and lambs at 1 mL/10Kg body weight.

Withholding Periods

Meat : 14 days
Milk : Do not use in dairy breeds of sheep.
Wool : N/A

Trade Advice

Export Slaughter interval: 28 days.

Accessories

This product can be used with any oral applicator designed to oral products to sheep. Contact your local reseller for the different options.

Body Weight (kg)	Dose Volume (mL)
11 - 15	1.5
16 - 20	2
21 - 30	3
31 - 40	4
41 - 50	5
51 - 60	6
61 - 70	7
71 - 80	8

