

SF00-SF06-0003

RANGER

Abamectin Plus Selenium Broad Spectrum Antiparasitic Drench for Sheep and Lambs



Active Content & Action

Abamectin 0.8 g/L, Selenium 0.4 g/L. Abamectin is part of the Macrocylic Lactones group of chemicals. It works by interrupting the nervous systems of parasites, leading to paralysis and eventual death. Selenium is a nutritional supplement provided to prevent white muscle disease.

Use

For the treatment and control of Abamectin sensitive strains of internal parasites in sheep (including Benzimidazole, Levamisole and Morantel resistant strains), nasal bot and itch mite of sheep and as an aid in the control of selenium deficiency in sheep including:

Adult and Immature Gastrointestinal Roundworms:

<i>Haemonchus contortus</i>	- Barber's pole worm (including inhibited L4 stage)
<i>Haemonchus placei</i>	- Large stomach worm
<i>Ostertagia circumcincta</i>	- Small brown stomach worm (including inhibited L4 stage)
<i>Trichostrongylus axei</i>	- Stomach hair worm
<i>Trichostrongylus colubriformis</i> , <i>Trichostrongylus vitrinus</i>	- Black scour worm
<i>Cooperia curticei</i> , <i>Cooperia oncophora</i>	- Small intestinal worm
<i>Nematodirus spathiger</i> , <i>Nematodirus filicollis</i>	- Thin necked intestinal worm
<i>Chabertia ovina</i>	- Large mouthed bowel worm
<i>Oesophagostomum columbianum</i>	- Nodule worm
<i>Oesophagostomum venulosum</i>	- Large bowel worm
<i>Trichuris ovis</i>	- Whipworm
<i>Strongyloides papillosus</i>	- Intestinal threadworm

Also Treats and Controls:

<i>Dictyoacaulus viviparus</i>	- Large lungworm
<i>Psorergates ovis</i>	- Itch mite
<i>Oestrus ovis</i>	- Nasal bot

Indications

Can be applied to all sheep. Do not use for sheep which are producing or may in the future produce milk products for human consumption.

Information About Selenium:

Ranger contains selenium. Excess selenium can cause toxicity issues in sheep, particularly if they are suffering liver damage from eating particular weeds. Where sheep are receiving sufficient selenium in their diet, it may be better to use an unselenised drench.

Period of Protection

This is a short acting product and can only be expected to work where resistance is not prevalent. Always conduct drench resistance testing and seek veterinary advice if you are concerned for livestock wellbeing.

Doses

1600 doses per 20L pack (average weight 50kg).

Pack Size(s) Available

20L

Australian Poisons Schedule

S5

Presentation

40 drums to a pallet

APVMA Approval No.:

80860



Dosage and Administration FOR EXTERNAL USE ONLY.

Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs). Do not underdose.

A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Use 1mL per 4Kg bodyweight.

Withholding Periods

Meat : 14 Days
Milk : Dispose of all milk collected from animals treated with this product
Woo : N/A

Trade Advice

Export Slaughter interval: 28 Days

Accessories

This product can be used with any oral applicator designed to apply drench products to sheep. Contact your local reseller for the different options.

11 - 16 kg	4.0 mL
17 - 20 kg	5.0 mL
21 - 30 kg	7.5 mL
31 - 40 kg	10.0 mL
41 - 50 kg	12.5 mL
51 - 60 kg	15.0 mL
61 - 70 kg	17.5 mL
71 - 80 kg	20.0 mL

